

ORIGINAL ARTICLE

PEER REVIEWED | OPEN ACCESS

Study of risk factors in patients hospitalized for psychotic disorders in the Dalal Xel mental health center in Fatick (Senegal)

Ibrahima Seck, Jean Augustin Diegane Tine, Abou SY, Matar BA, Khadim Niang, François Ngor Faye, Anta Tal-Dia

ABSTRACT

Aims: Psychiatric disorders are characterized by the complexity of their psychodynamic mechanisms and the inadequacy of therapeutic means. This study aims to study the risk factors of psychotic pathologies hospitalized in Dalal Xel of Fatick. **Methods:** This is a cross-sectional, retrospective, registrable, descriptive and analytical study carried out between 02 June and 21 July 2014 in the Dalal Xel mental health center in Fatick. All patients hospitalized in this facility were included between November 2003 and April 2014 and their medical records were retrieved. Simple logistic regression was used. **Results:** Of the 7274 patient's record collected, the average age was 34.32 years (± 20.32), men were the majority at 61.2% and 57.6% were from rural areas. Serer represented 39.6% of cases. The 34.9% were out of school. Psychotic forms were mostly represented (56.8%). The main factors related to psychosis after adjustment

are age between 16 and 40 years (OR = 8.2 [6.6–10.3]), marriage (OR = 0.7 [0.6–0.9]), the death of the father (OR = 1.3 [1.1–1.5]), the death of the mother (OR = 1.4 [1.1–1.9]). **Conclusion:** Psychotic disorders are influenced by the socio-family experience of the individual. Marriage, orphan status and family support have been identified as risk factors in psychosis, hence the importance of involving the family in patient follow-up and in the development of mental health policies.

Keywords: Dalal Xel, Fatick, Psychiatric pathologies, Risk factors, Senegal

How to cite this article

Seck I, Tine Jad, SY A, Matar BA, Niang K, Faye FN, Tal-Dia A. Study of risk factors in patients hospitalized for psychotic disorders in the Dalal Xel mental health center in Fatick (Senegal). *Edorium J Public Health* 2017;4:20–26.

Article ID: 100011P16IS2017

doi:10.5348/P16-2017-11-OA-3

Ibrahima Seck¹, Jean Augustin Diegane Tine², Abou SY³, Matar BA³, Khadim Niang², François Ngor Faye⁴, Anta Tal-Dia⁵

Affiliations: ¹Professor, Director of studies, Public Health and Preventive Medicine, ISED, Dakar, Senegal; ²Assistant, Public Health and Preventive Medicine, ISED, Dakar, Senegal; ³Physician, Hospital Fann, Psychiatry, Dakar, Senegal; ⁴Director, Dalal Xel Mental Health Center, Dakar, Senegal; ⁵Professor, Director of ISED, Public Health and Preventive Medicine, ISED, Dakar, Senegal.

Corresponding Author: Ibrahima Seck, Institut de Santé et Développement (ISED), Dakar-Fann, Senegal, BP 16 390; Email: ibouseck304@gmail.com

Received: 02 January 2017

Accepted: 07 March 2017

Published: 27 March 2017

INTRODUCTION

Psychotic pathologies have the particularity of being often chronic and costly. According to Charrier, the direct costs associated with schizophrenia range from US \$16 billion in the United States to US \$204 billion in the United Kingdom and \$79 billion in Canada [1]. Developing countries, particularly those in Africa, are no exception. Those who are already struggling with their inadequate health systems and on a small budget do not

have the means to cope with the growing mental health problems [2].

Since 1956, Senegal has setup psychiatric structures to provide a sanitary environment for the mentally ill with the creation of the neuropsychiatry department of the Fann hospital. In 1994, the first private non-profit psychiatric structure was created as the mental health center “Dalal Xel” in Thiès. Then in 2003, another health center “Dalal Xel” was opened in Fatick. This partnership between the Order of “Saint-Jean de Dieu” and the State of Senegal has increased the coverage and quality of mental health services. Dalal Xel’s mental health care providers offer a range of preventive and curative services to the population. [3].

This study investigated the risk factors of patients hospitalized for psychotic disorders in the Dalal Xel mental health center in Fatick, Senegal. The specific objectives were:

- describe the socio-demographic characteristics (individual and family) of patients hospitalized in the Dalal Xel mental health center in Fatick
- describe the occupational characteristics of patients hospitalized at Dalal Xel mental health center in Fatick
- determine the frequencies of the various pathologies hospitalized in Dalal Xel mental health center in Fatick
- analyze socio-demographic factors related to psychotic illnesses hospitalized in Dalal Xel mental health center in Fatick
- analyze the occupational factors related to the psychotic pathologies hospitalized in the Dalal Xel mental health center of Fatick

MATERIALS AND METHODS

Presentation of Dalal Xel health center

Dalal Xel health center is a health structure of the Catholic Church owned by the Hospitaller Order of the brothers of Saint John of God located in the region of Thiès. The Hospitaller Order of the brothers of Saint John of God was founded in 1550 by Juan Ciudad who became John of God (1495–1550).

In Senegal, the brothers of Saint John of God created the hospital of Saint Jean de Dieu in Thiès in 1984. In view of the impact of mental illness on the Senegalese population, they decided to create the mental health center of Senegal. Dalal Xel of Thiès in 2000.

The mental health center of Fatick is located at the entrance of the town of Fatick on National No. 1 150 km from Dakar and 65 km from Mbour.

The Dalal Xel system is under the guidance of the Consultative Council (CAC).

Each Dalal Xel is managed by a steering committee composed of the head of the health center, the chief

medical officer and an administrative and financial manager.

The Dalal Xel health center in Fatick is subdivided into four departments: management, outpatient ward, hospital wards and occupational therapy.

The Dalal Xel mental health center in Fatick has a small staff. By September 2014, there were 10 caregivers including two psychiatrists, one psychopathology nurse and seven nursing assistants.

The activities of the psychiatric center are general psychiatric consultation, adult hospitalization and occupational therapy and occupational therapies such as gardening, weaving, games of mind (card game, lady’s play), etc.

Type of study

This was a cross-sectional, retrospective, descriptive and analytical study.

Study period

The quantitative study covered the period from 02 June to 21 July 2014.

Study population and sampling

This study was exhaustive, on the basis of the hospitalization register and patient records. It was included all patients who were recorded in the hospitalization records of the structure between November 2003 and April 2014. It was excluded those registries that did not have a diagnosis clearly identified by the psychiatrist.

Procedure for collecting and entering data

In each hospitalization unit, there is a register at the nurses’ room and a file at the doctor’s office.

All the patient files, found in the hospitalization units of the structure and belonging to the study period, were collated in a seizure mask made on Microsoft Access 2007 by two seizure operators trained and taken care of by the structure.

Data analysis

The analysis of the data was done with Epi-info 3.5.3 and with software R 3.1.1. The qualitative variables were described with the frequency and 95% confidence interval and the quantitative variables by mean, standard deviation and quantiles. Tables and charts were built on Microsoft Excel 2007.

Variables have been crossed to meet the expectations of the determinants objectives. This was done using the tests of KHI2 (comparison of two proportions) or

Student (comparison of two averages) according to their applicability conditions with a significance threshold of $p = 0.05$. The odds ratio, surrounded by its confidence interval, quantified the strength of the link.

A multivariate analysis was performed with a simple logistic regression for the determination of factors related to psychosis, with an adjustment to the variables related to the individual, family and occupational characteristics while ensuring the conditions of applicability with a threshold of significance set at 5%. Variables with p values less than 0.25 in bivariate analysis were used for modeling [4].

The lrtest (likelihood ratio) and AIC allowed the choice of the last one with the step-by-step descending method. Finally, the Hosmer Lemeshow test made it possible to test the suitability of the final model. [5]

In the models, the associations were measured by the odds ratio with their confidence intervals.

Ethical considerations

This study was initiated as part of an evaluation of the activities of the Dalal Xel mental health center in collaboration with the head of the health center. The data is collected anonymously. No judgment of subjective, personified or group value that may taint the personality of the individuals and the institution will be published in this study. No remuneration or financial or material compensation was offered to the patients.

RESULTS

Clinical features

Between November 2003 and April 2014, there were 7274 patients who were hospitalized at the Dalal Xel psychiatric center in Fatick. By filing the records, it was found that 91.55% (6660 patients) of the cases had a diagnosis of discharge. Of these patients with a diagnosis, 72.8% (4848 cases) had psychiatric illnesses, 2.5% (1766 cases) had neurological diseases as a cause of hospitalization, and the rest had general medicines such as gastric disturbances, infectious syndromes, etc.

In the psychiatric illness group, 77.6% (3763 cases) had psychoses, 15% (728 cases) had mood disorders (depression, mania, bipolar disorder) and 7.4% (357 cases) were neurotic according to the psychiatric classification of the French school.

In the psychotic group, using ICD10, we found that 34.3% (1291 cases) had schizophrenic disorders, 32.1% (1207 cases) had acute delusional disorders, 15% (566 cases) had mental disorders related to psychoactive substances. Chronic hallucinatory psychotic disorders and non-dissociative psychotic disorders accounted for 8.9% (336 cases) and 6.9% (225 cases, respectively). The mental disorders associated with puerperality were

minority at 2.8% (108 cases). Table 1 summarizes the distribution of cases according to ICD 10.

Socio-demographic characteristics

The mean age of hospitalized patients was 34.3 years with a standard deviation of 20.3 years. Men were in the majority with a sex ratio (M / F) of 1.5. The unmarried, the childless, the uneducated were more frequent. Most of the patients were orphans. Table 2 summarizes the socio-demographic characteristics of the patients.

Professional features

The majority of patients reported having a vocational qualification. The 93.2% had a job in the private sector. The 36.3% had a domestic job. It had 1.4% of learners and 2.9% of unemployed. Table 3 summarizes the occupational characteristics of the patients.

Analytical section

Multivariate analysis has shown that age, educational level, orphan status, family support, geographic origin are risk factors for psychosis. Table 4 gives the results of the multivariate analysis made by a simple logistic regression.

DISCUSSION

This retrospective study was confronted with missing values in the exploration of hospitalization records. Consultation activities were not explored because of the

Table 1: Distribution of cases of psychotic disorders according to ICD 10

Diagnoses according to ICD 10	Absolute Frequencies (n)	Relative Frequencies (%)
Schizophrenic disorders	1291	34,3
Acute delusional disorders	1207	32,1
Mental disorders related to psychoactive substances	566	15,0
Chronic hallucinatory psychotic disorders	336	8,9
Chronic Non-dissociative psychotic disorders	255	6,9
Mental disorders associated with puerperality	108	2,8
Total	3763	100,0

Table 2: Distribution of hospitalized patients by socio-demographic

Socio-demographic characteristics	n (%) or μ ($\pm\epsilon$)
Age (year)	34,3 (\pm 20.3)
Gender: Male	4435 (61.2)
Marital Status	
Single	4112 (57.1)
Married	2453 (34.1)
Divorced	373 (5.2)
Widower	259 (3.6)
Patient's number of children	
No Child	6671 (91.8)
Between 1 to 6 children	503 (6.9)
More than 6 children	95 (1.3)
Religion	
Muslim	6751 (92.9)
Catholic	516 (7.1)
Level of study	
University	295 (4.1)
Secondary school	1298 (18.1)
Primary	1917 (26.7)
Quranic learned only	1169 (16.3)
No school education	2509 (34.8)
Geographic origin	
Urban	2970 (42.4)
Rural	4039 (57.6)
Marital status of the family of origin	
Polygamous	4054 (59.0)
Monogamous	2812 (40.9)
Divorced	1 (0.01)
Parents experience	
Parents in life	3303 (51.0)
Deceased Father	1760 (27.2)
Deceased Mother	375 (5.8)
Deceased Parents	1038 (16.0)
Good family support (Yes)	6698 (97.5)

incompleteness of socio-demographic information and especially clinical diagnosis. But we had a statistically representative number of records to do the logistic regression.

A psychosis is a mental disorder characterized by the loss of contact with reality, a disorganization of the personality, and the delirious transformation of experience. Psychoses are thus distinguished from

Table 3: Distribution of hospitalized patients by occupational characteristics

Professional Features	n (%) or μ ($\pm\epsilon$)
Existence of professional qualification (yes)	4690 (68.3)
Employment sector	
Public	292 (6.8)
Private/ liberal	3995 (93.2)
Type of occupation	
Housework	1555 (36.3)
Farmer/ Breeder	985 (23.0)
Trader	494 (11.5)
Worker/ artisan	459 (10.7)
Teacher	179 (4.2)
Esthetician/ tailor	169 (3.9)
Driver	162 (3.8)
Fisher	62 (1.4)
Artist	56 (1.3)
Restaurant/hotel	50 (1.2)
Administrative	46 (1.1)
Security/ military	36 (0.8)
Health worker	16 (0.4)
Contractor	15 (0.3)
Religious	3 (0.1)
Unemployed (yes)	135 (2.9)
Learner status (student/school boy (yes)	99 (1.4)
Retired status (yes)	249 (10)

neuroses, in which the patient retains the notion of the real, even if sometimes he does not understand the interpretation. In a neurosis, the subject retains a critical mind towards his illness. Chronic psychoses include chronic delusions, schizophrenia, paranoia, chronic hallucinatory psychosis (psychiatric hallucinations). Psychotic disorders, opening on a delusional experience, occur in several circumstances [6].

Psychoses are very serious disease because of their physical, economic and social consequences. This study estimated the extent of these psychotic pathologies to be 3763 inpatients between November 2003 and April 2014 in this small 23-bed mental health center. On an average we found one case of psychotic hospitalization per day for more than 10 years.

The proportional morbidity of psychoses in hospitalization in this structure is 77.6%. It is thus the leading cause of hospitalization, which is explained by its seriousness. The Kirkbride meta-analysis showed a lower prevalence of psychotic disorders (4‰) but was in the general population [7]. Schizophrenic disorders accounted for 34.3% of cases of psychosis.

Table 4: Psychotic risk factors by logistic regression

	Psychotic		
	P	ORa [IC 95%]	ORbrut [IC 95%]
Age			
1–15 years	ref	-	-
16–40 years	2×10 ⁻¹⁶	6.5 [5.0–8.5]	8.2 [6.6–10.3]
>40 years	2.2×10 ⁻¹⁶	5.69 [4.1–7.7]	5.2 [4.1–6.6]
Patient marital status			
Single	ref	-	-
Married	1.3×10 ⁻⁵	0.7 [0.6–0.8]	0.8 [0.7–0.9]
Divorced	0.2	-	1.3 [1.04–1.6]
Widow	7.22×10 ⁻⁷	0.4 [0.2–0.5]	0.3 [0.2–0.5]
Level of education			
No education	ref	-	-
University	0.001	1.6 [1.3–2.3]	1.8 [1.4–2.3]
Secondary	1.5×10 ⁻¹¹	1.9 [1.5–2.3]	2.0 [1.7–2.3]
Primairy	4.8×10 ⁻⁸	1.5 [1.3–1.8]	1.6 [1.4–1.9]
Coranic teaching	0.08	-	1.4 [1.2–1.6]
Childhood environment			
Urban	ref	-	-
Rural	0.08	-	0.80 [0.7–0.88]
Parental status			
Parents in life	ref	-	-
Deceased father	0.0001	1.3 [1.1–1.5]	0.8 [0.7–0.9]
Deceased mother	0.006	1.4 [1.1–1.9]	1.5 [1.2–1.9]
Deceased Parents	0.22	-	1.5 [1.3–1.7]
Good family support			
No	ref	-	-
Yes	4.5×10 ⁻¹²	0.2 [0.1–0.3]	0.20 [0.1–0.29]
Existence of one qualification			
Yes	ref	-	-
No	4.0×10 ⁻¹²	1.5 [1.3–1.8]	1.3 [1.2–1.5]
Learner status			
No	ref	-	-
Yes	0.002	0.3 [0.1–0.6]	0.5 [0.3–0.9]

†ref: References range

A rural geographic origin protected against psychosis (OR = 0.8 [0.7–0.88]) [8–10]. Urbanization would be a factor in aggravating or decompensating psychosis with a very high risk [8, 11–13]. Lifestyle and education

different from the city to the countryside. In Senegal, rural life is very communal with broad social support that differs from the urban organization that is becoming more and more nuclear. With the difficulties of life in the countryside, education is based on traditional, religious and resilience bases called in wolof moun.

Exposure to psychosis is higher among people aged 16 to 40 years. This result agrees with the age of schizophrenia [9, 14–18]. This age group represents a large part of the working population, hence the importance of the impact of psychiatric pathologies on the development of a country [19–21].

The married couple are less exposed to psychosis (OR = 0.7 [0.6–0.8] or psychosis which is a chronic disease, usually dyssocial does not allow the patient to be able to marry. The orphan is exposed to psychosis through the lack of a solid social fabric and a parental framework in the psychological development of the person, in medical assistance and therapeutic follow-up.

This lack of follow-up of certain orphans may trigger predisposing factors or the complication of a psychotic state that may associate it with mood disorders. This result agrees with the usefulness of family support in the medical follow-up of the psychotic (OR = 0.2 [0.1–0.3]). Therapists often involve the family in the follow of the sick at home. They understood the importance of the family in the follow-up of psychotics [10, 11, 19, 20]. To facilitate the involvement of families, the accessibility of mental health services should be improved through decentralization of mental health care to different levels of the health pyramid by identifying the appropriate service packages [12, 22, 23, 24].

The risk of psychosis is higher in patients without qualifications (OR = 1.5 [1.3–1.8]). Student status would protect against psychosis with OR = 0.3 [0.1–0.6]. This result would also be related to the mental handicap of the disease, which means that patients do not have the opportunity to follow the normal educational cycle, which leads to an early termination of studies and a lack of qualification [13, 25].

CONCLUSION

Fatick’s Dalal Xel Mental Health Center is a small, 23-bed facility that receives an average of 700 patients per year. It hosts all psychiatric pathologies, neurological pathologies and even medical emergencies.

The risk factors for psychotic disorders are age, marital status, orphan status and inadequate family support.

Acknowledgements

We acknowledge al Dal Xel center and Fann Hospital Psychiatric Staff who help us in the work.

Author Contributions

Ibrahima Seck – Substantial contributions to conception and design, Revising it critically for important intellectual content, Final approval of the version to be published

Jean Augustin Diegane Tine – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Final approval of the version to be published

Abou SY – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Matar BA – Acquisition of data, Analysis and interpretation of data, Revising it critically for important intellectual content, Final approval of the version to be published

Khadim Niang – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Final approval of the version to be published

François Ngor Faye – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Anta Tal-Dia – Substantial contributions to conception and design, Revising it critically for important intellectual content, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2017 Ibrahima Seck et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

- World economic forum & harvard School of public health. The global economic burden of non-communicable diseases. Genève: Forum économique mondial; 2001. [Available at: http://www3.weforum.org/docs/WEF_Harvard_HE_Global_Economic_Burden_Non_Communicable_Diseases_2011.pdf]
- Cohidon C. Prevalence of mental health disorders and consequences on occupational activity in France in the mental health in the general population: Images and realities survey. France: Institute for Health Surveillance; 2007.
- Tine Benoit Addiction: Sociological Study of a Form of Deviance in Senegal. Paris: Publibook; 2008.
- Collet D. Modeling binary data. 2ed. London, New York: Chapman and Hall/CRC; 2003. p. 387.
- Hosmer DW, Lemeshow S. Applied logistic regression. New York: John Wiley; 1989. p. 307.
- Besançon G. Manual of Psychopathology-Psychoses. Paris: Dunod; 1993.
- Kirkbride JB, Errazuriz A, Croudace TJ, et al. Systematic review of the incidence and prevalence of schizophrenia and other psychoses in England. Conducted for the Department of Health Policy Research Programme. 2012. [Available at: <http://www.psychiatry.cam.ac.uk/files/2014/05/Final-report-v1.05-Jan-12.pdf>]
- Harrison G, Fouskakis D, Rasmussen F, Tynelius P, Sipos A, Gunnell D. Association between psychotic disorder and urban place of birth is not mediated by obstetric complications or childhood socio-economic position: A cohort study. *Psychol Med* 2003 May;33(4):723–31.
- Jones PB. Schizophrenia & psychotic disorders: Current epidemiological highlights. *CEPiP* 2011;1(3)Indd 6.
- Friesen B, Katz-Leavy J, Nicholson J. Supporting parents With Mental Health Needs in Systems of Care. Washington, DC: Technical Assistance Partnership for Child and Family Mental Health; 2011. [Available at: http://www.air.org/sites/default/files/downloads/report/Supporting_Parents_With_Mental_Health_Needs_2_0.pdf]
- Leclerc L. Being a caregiver close to a person living with a mental health disorder in a high-speed society. *The partner Quebec* 2012;20(4):6.
- Diop B, Collignon R, Gueye M. Presentation of the collaborative study of the OMS. On strategies for extending mental health care. *Psychopath* 1976;(12):173–88.
- Perälä J, Suvisaari J, Saarni SI, et al. Lifetime prevalence of psychotic and bipolar I disorders in a general population. *Arch Gen Psychiatry* 2007 Jan;64(1):19–28.
- Allardyce J, Boydell J, Van Os J, et al. Comparison of the incidence of schizophrenia in rural Dumfries and Galloway and urban Camberwell. *Br J Psychiatry* 2001 Oct;179:335–9.
- Bamrah JS, Freeman HL, Goldberg DP. Epidemiology of schizophrenia in Salford, 1974–84. Changes in an urban community over ten years. *Br J Psychiatry* 1991 Dec;159:802–10.
- Baudis P. A comparison of first admissions of mania in Prague, Aarhus and London:

- A transcultural study linked up with the international pilot study of schizophrenia of WHO. *Social Psychiatry* 1977;12(4):185–93.
17. Bhugra D, Leff J, Mallett R, Der G, Corridan B, Rudge S. Incidence and outcome of schizophrenia in whites, African-Caribbeans and Asians in London. *Psychol Med* 1997 Jul;27(4):791–8.
 18. Boydell J, Van Os J, Lambri M, et al. Incidence of schizophrenia in south-east London between 1965 and 1997. *Br J Psychiatry* 2003 Jan;182:45–9.
 19. de Alarcon J, Seagroatt V, Sellar C, Goldacre M. Population-based trends in treatment rates in psychiatry in Oxfordshire, 1975–1986. *J Public Health Med* 1993 Mar;15(1):93–102.
 20. Fearon P, Kirkbride JB, Morgan C, et al. Incidence of schizophrenia and other psychoses in ethnic minority groups: Results from the MRC AESOP Study. *Psychol Med* 2006 Nov;36(11):1541–50.
 21. Gater R, Amaddeo F, Tansella M, Jackson G, Goldberg D. A comparison of community-based care for schizophrenia in south Verona and south Manchester. *Br J Psychiatry* 1995 Mar;166(3):344–52.
 22. Kirkbride JB, Fearon P, Morgan C, et al. Heterogeneity in incidence rates of schizophrenia and other psychotic syndromes: Findings from the 3-center AeSOP study. *Arch Gen Psychiatry* 2006 Mar;63(3):250–8.
 23. Kirkbride JB, Fearon P, Morgan C, et al. Neighbourhood variation in the incidence of psychotic disorders in Southeast London. *Soc Psychiatry Psychiatr Epidemiol* 2007 Jun;42(6):438–45.
 24. Kirkbride JB, Morgan C, Fearon P, Dazzan P, Murray RM, Jones PB. Neighbourhood-level effects on psychoses: Re-examining the role of context. *Psychol Med* 2007 Oct;37(10):1413–25.
 25. Morgan C, Dazzan P, Morgan K, et al. First episode psychosis and ethnicity: Initial findings from the AESOP study. *World Psychiatry* 2006 Feb;5(1):40–6.

Access full text article on
other devices



Access PDF of article on
other devices

